INDEX

Page

ABSORPTION OF COPPER FROM THE SOIL BY POTATO PLANTS..... 281-287
Acer—
rubrum, host of *Xylotrechus colonus*..... 195-198
saccharum, organic acids..... 221-229
Achillea ptarmica, susceptibility to *Puccinia triticina*..... 152-172
Acid—
amino, in Georgia velvet beans..... 15
citric, in grapefruit..... 263-279
diamino, in Georgia velvet bean..... 15
hydrocyanic, in Sudan grass..... 125-138
malic, in *Pyrus coronaria*..... 221-229
phosphoric, effect on hydrocyanic acid in Sudan grass..... 135
succinic, in *Pyrus coronaria*..... 221-229
sulphuric, effect on hydrocyanic acid in Sudan grass..... 133-135
Acid producers in corn meal..... 181-188
Acid-sugar ratio in grapefruit..... 263-279
Acids—
effect on hydrocyanic acid in Sudan grass. 133-135
organic, of *Pyrus coronaria*, *Rhus glabra*, and *Acer saccharum*..... 221-229
Aconitum—
Fischeri, resistance to teliospores of *Puccinia triticina*..... 155-172
Lycoctonum, susceptibility to *Puccinia triticina*..... 152-172
Napellus, susceptibility to *Puccinia triticina*..... 152-172
Actaea spicata, resistance to teliospores of *Puccinia triticina*..... 155-172
Actinomyces sp. in corn meal..... 187-188
AECIAL STAGE OF THE ORANGE LEAFRUST OF WHEAT, PUCCINIA TRITICINA ERIKS..... 151-172
Aecidium—
Anchusae, pathogenicity on wheat..... 152
Berberidis, in life cycle of *Puccinia graminis*..... 151-172
Ranunculacearum γ *Thalictri flavi*. Syn. *Aecidium Thalictri flavi*.
Thalictri flavi, rust on *Agropyron repens*... 164
Aegopodium Podagraria, susceptibility to *Puccinia triticina*..... 152-172
Agropyron—
caninum, resistance to aeciospores of *Puccinia triticina*..... 163-172
cristatum, resistance to aeciospores of *Puccinia triticina*..... 163-172
desertorum, resistance to aeciospores of *Puccinia triticina*..... 163-172
inerme, resistance to aeciospores of *Puccinia triticina*..... 163-172

Page

Agropyron—Continued.
repens, resistance to aeciospores of *Puccinia triticina*..... 163-172
Richardsonii, host of *Puccinia Agropyri*.... 165
Smithii, resistance to aeciospores of *Puccinia Agropyri*..... 165
tenerum, resistance to aeciospores of *Puccinia triticini*..... 163-172
Agrostemma Githago, susceptibility to *Puccinia triticina*..... 152-172
Agrostis—
borealis, host of *Puccinia triticina*..... 164-172
orthogonia. Syn. *Porosagrotis orthogonia*.
Akis muricata. Syn. *Embaphion muricatum*.
Albumen, egg, effect on growing chicks..... 146
Albumin in Georgia velvet bean..... 15
Alfalfa, effect of soil temperature on development of nodules..... 17-31
Alkaline solution, effect on hydrocyanic acid in Sudan grass..... 135-136
Alternaria sp. on pitted grapefruit..... 277
Ambrosia artemisiifolia, glucosid..... 4
Amino acid. See Acid, amino.
Ammodramus savannarum bimaculatus, enemy of *Porosagrotis orthogonia*..... 313
Amygdalus persica—
effect of temperature on damage from rots 452-465
host of *Lecanium corni*..... 191
Anchusa—
arvensis, host of aecial stage of *Puccinia rubigo-vera*..... 151-172
officinalis, host of aecial stage of *Puccinia rubigo-vera*..... 151-172
Ancylostoma—
caninum, nonhemolytic effect..... 383
duodenale, hemotoxins..... 382-432
Anemone—
canadensis, resistance to teliospores of *Puccinia triticina*..... 155-172
cylindrica, resistance to teliospores *Puccinia triticina*..... 155-172
japonica, resistance to *Puccinia triticina*.. 155-172
ranunculoides, susceptibility to *Puccinia triticina*..... 152-172
Anoplocephala—
perfoliata, hemotoxins not found..... 382
plicata, hemotoxins not found..... 382
Anthocyanin in maize..... 2-31
Anthoxanthum odoratum, host of *Puccinia triticina*..... 164-172
Anticoagulins in hookworms..... 418-420
Aphids, carriers of mosaic of Chinese cabbage..... 173-178
Aphis, sugar-beet root. See *Pemphigus betae*.
Apple blotch. See *Phyllosticta solitaria*.

Page

Apple, crab. See *Pyrus coronaria.*
Aquilegia—
alpina, resistance to teliospores of *Puccinia triticina* ... 155–172
canadensis, resistance to teliospores of *Puccinia triticina* ... 155–172
chrysantha, resistance to teliospores of *Puccinia triticina* ... 155–172
glandulosa, resistance to teliospores of *Puccinia triticina* ... 155–172
olympica, resistance to teliospores of *Puccinia triticina* ... 157–172
Skinneri, resistance to teliospores of *Puccinia triticina* ... 155–172
vulgaris, resistance to teliospores of *Puccinia triticina* ... 155–172
Army cutworm. See *Chorizagrotis auxiliaris.*
Arrhenatherum elatis, resistance to aeciospores of *Puccinia triticina* ... 163–172
Ascaris—
conocephala, cause of anemia ... 385–432
lumbricoides, hemotoxins ... 385–432
Ash, bone, effect on growing chicks ... 145–149
Ash. See *Fraxinus* sp.
ASH CONTENT OF THE AWN, RACHIS, PALEA, AND KERNEL OF BARLEY DURING GROWTH AND MATURATION ... 433–449
Aspergillus—
candidus in corn meal ... 187–188
flavus in corn meal ... 181–188
fumigatus in corn meal ... 185–188
niger in corn meal ... 181–188
ochraceous in corn meal ... 187–188
repens in corn meal ... 181–188
tamari in corn meal ... 185–188
terreus in corn meal ... 185–188
Awn of barley, ash content ... 433–449
Bacillus—
coli communis, effect on aspartic acid ... 224
niger in corn meal ... 186–188
phytophthorus, cause of balckleg potato tuber-rot ... 81–92
radicola—
effect of sulphur ... 103–110
used in tests of effect of soil temperature on nodule development ... 20–31
tetanus, blood toxin ... 381
Bacteria—
in cheese ... 93, 100
in corn meal ... 181–188
Bacterium aerogenes in corn meal ... 183–188
Barbaraea vulgaris, susceptibility to *Puccinia triticina* ... 152–172
Barley, ash content ... 433–449
Bartlett, H. H., and Sando, Charles E.:
NOTES ON THE ORGANIC ACIDS OF PYRUS CORONARIA, RHUS GLABRA, AND ACER SACCHARUM ... 221–229
OCCURRENCE OF QUERCETIN IN EMERSON'S BROWN-HUSKED TYPE OF MAIZE ... 1–31
Bean, velvet. See *Stizolobium deeringianum.*
Beetle, pine. See *Dendroctonus monticolae.*

Page

Beetles, cerambycid, host-selection principle ... 189–220
Beets, sugar, dryrot canker ... 47–52
Berberis vulgaris—
failure to become infected from basidiospores of *Puccinia rubigo-vera* ... 152–172
susceptibility to *Puccinia triticina* ... 152–172
BIOLOGICAL ANALYSIS OF THE SEED OF THE GEORGIA VELVET BEAN, STIZOLOBIUM DEERINGIANUM ... 5–15
BIOLOGY OF EMBAPHION MURICATUM ... 323–334
Black locust. See *Robinia pseudacacia.*
Black mold rot, caused by *Rhizopus nigricans* ... 451–465
BLACKLEG POTATO TUBER-ROT UNDER IRRIGATION ... 81–92
Blight, chestnut. See *Endothea parasitica.*
Blotch—
apple. See *Phyllosticta solitaria.*
plum. See *Phyllosticta congesta.*
Blue mold rot. See *Penicillium expansum.*
Bone ash, effect on growing chicks ... 145–149
Bonnetia compta, parasite of *Porosagrotis orthogonia* ... 313
Boshnakian, Sarkis, and Leighty, Clyde E.:
GENETIC BEHAVIOR OF THE SPELT FORM IN CROSSES BETWEEN TRITICUM SPELTA AND TRITICUM SATIVUM ... 335–364
Böving, Adam G., and Wade, J. S.: BIOLOGY OF EMBAPHION MURICATUM ... 323–334
Brassica—
japonica, mosaic disease ... 173–178
pekinensis, mosaic disease ... 173–178
rapa, mosaic disease ... 173–178
Bromus—
ciliatus, host of *Puccinia Agropyri* ... 165
Porteri, host of *Puccinia alternans* ... 165
Brooks, Charles, and Cooley, J. S.:
TEMPERATURE RELATIONS OF STONE FRUIT FUNGI ... 451–465
TRANSPORTATION ROTS OF STONE FRUITS AS INFLUENCED BY ORCHARD SPRAYING ... 467–477
Brown-husked maize, quercetin ... 1–31
Brown rot. See *Sclerotinia cinerea.*
Bustomum phlebotomum, hemptoxins ... 415–432
Cabbage, Chinese. See *Brassica pekinensis.*
Calcium—
carbonate, effect on—
growing chicks ... 139–149
plant growth ... 102–110
toxicity of copper sulphate ... 281
lactate, effect on growing chicks ... 145–149
sulphate, effect on plant growth ... 102–110
Callidium—
antennatum, host selection ... 194–220
janthinum, host selection ... 203–220
Camassia esculenta, resistance to teliospores of *Puccinia triticina* ... 155–172
Campanula rotundifolia, susceptibility to *Puccinia triticina* ... 152–172

Page

Canker of sugar beets 47-52
Calvin, J. W., et al.: NUTRIENT REQUIREMENTS OF GROWING CHICKS: NUTRITIVE DEFICIENCIES OF CORN 139-149
Camallanus americanus, oxyhemoglobin in body fluid 392-432
Carbonate—
calcium, effect on—
growing chicks 139-149
plant growth 102-110
toxicity of copper sulphate 281-287
sodium, effect on hydrocyanic acid in Sudan grass 135-136
Casein, effect on growing chicks 139-149
Castanea dentata, host of—
Hyperplatys maculatus 217-219
Xylotrechus colonus 195-198
Cattle hookworm. See *Bustomum phlebotomum*.
Cattle, maintenance requirement 115-121
Celtis occidentalis, host of—
Cyllene pictus 198-203
Molorchus bimaculatus 212-213
Centaurea Cyanus, susceptibility to *Puccinia triticina* 152-172
Cerambycid beetles, host-selection principle 189-220
Cercis canadensis, immunity to *Molorchus bimaculatus* 212-213
Cestode hemolysins 420-432
Cheese, bacteria 93-100
Cherries, effect of—
orchard spraying on transportation rots .. 467-477
temperature on damage from rots 441-465
Chestnut. See *Castanea dentata*.
Chestnut blight. See *Endothea parasitica*.
Chinese cabbage. See *Brassica pekinensis*.
Chlorid, sodium, effect on growing chicks .. 145-149
Chorizagrotis auxiliaris, comparative destructiveness of *Porosagrotis orthogonia* 289
Cimicifuga racemosa, resistance to teliospores of *Puccinia triticina* 155-172
Citellus richardsoni, enemy of *Porosagrotis orthogonia* 313
Citric acid. See Acid, citric.
Citromyces sp. in corn meal 187-188
Cladosporium sp. on pitted grapefruit 277
Clematis—
Douglasii, resistance to teliospores of *Puccinia triticina* 155-172
flammula, probable aecial host of *Puccinia triticina* 153-172
Fremontii, resistance to teliospores of *Puccinia triticina* 155-172
heraclaefolia, resistance to teliospores *Puccinia triticina* 155-172
ligusticifolia, resistance to teliospores of *Puccinia triticina* 155-172
orientalis, resistance to teliospores of *Puccinia triticina* 155-172
paniculata, resistance to teliospores of *Puccinia triticina* 157-172
recta, resistance to teliospores of *Puccinia triticina* 155-172

Page

Clematis—Continued.
virginiana, resistance to teliospores of *Puccinia triticina* 155-172
vitalba, probable aecial host of *Puccinia triticina* 153-172
Clonorchis sinensis, hemotoxins 380-432
Colletotrichum sp. on pitted grapefruit 277
COMPARATIVE VIGOR OF F_1 WHEAT CROSSES AND THEIR PARENTS 53-63
COMPARISON OF THE PECTINASE PRODUCED BY DIFFERENT SPECIES OF RHIZOPUS, A 371-377
Coniferous tree seed, storage 479-510
Cook, F. C.: ABSORPTION OF COPPER FROM THE SOIL BY POTATO PLANTS 281-287
Cooley, J. S., and Brooks, Charles: TEMPERATURE RELATIONS OF STONE FRUIT FUNGI 451-465
Copper, absorption from soil by potato plants 281-287
Copper sulphate, toxicity to potato plants .. 281-287
Corn, nutritive defeciencies 139-149
Corn meal, flora 179-188
Cornus florida, host of—
Hyperplatys maculatus 217-219
Molorchus bimaculatus 212-213
Coronaria flosculi, susceptibility to *Puccinia triticina* 152-172
Corticium vagum, cause of dryrot canker of sugar beets 47-52
Crab apple. See *Pyrus coronaria*.
Craighead, F. C.: HOPKINS HOST-SELECTION PRINCIPLE AS RELATED TO CERTAIN CERMBYCID BEETLES 189-220
Cutworm—
army. *Chorizagrotis auxiliaris*.
pale western. See *Porosagrotis orthogonia* 289-322
Cyanidin, glucosid, in maize 2-31
Cyllene—
crinicornis, host selection 203
pictus, host selection 193-220
Delphinium—
ajacis, resistance to teliospores of *Puccinia triticina* 155-172
"Belladonna," resistance to teliospores of *Puccinia triticina* 155-172
consolida, resistance to teliospores of *Puccinia triticina* 155-172
geyeri, resistance to teliospores of *Puccinia triticina* 155-172
Dendroctonus monticolae, parasite on *Pinus contorta* 189-220
Dextrose in grapefruit 263-279
Diamino acid. See Acid, diamino.
Digestion, liberation of hydrocyanic acid 137
Diphyllobothrium latum, hemotoxins 381-432
Dipotassium phosphate, effect on growing chicks 145-149
Dipylidium caninum, hemotoxins 382-432
Dogwood. See *Cornus florida*.
DRYROT CANKER OF SUGAR BEETS, A 47-52

Page

Echium vulgare, susceptibility to *Puccinia triticina* ... 152–172
Edson, H. A., and Shapovalov, M.: BLACK-LEG POTATO TUBER-ROT UNDER IRRIGATION ... 81–92
EFFECT OF SOIL TEMPERATURE UPON THE DEVELOPMENT OF NODULES ON THE ROOTS OF CERTAIN LEGUMES ... 17–31
Egg albumen, effect on growing chicks ... 146
Eleodes conlusum, resemblance to *Embaphion muricatum* ... 323
Elymus—
australis, resistance to aeciospores *Puccinia triticina* ... 163–172
canadensis, resistance to aeciospores of *Puccinia triticina* ... 163–172
glaucus, resistance to aeciospores of *Puccinia triticina* ... 163–172
triticoides, resistance to aeciospores of *Puccinia triticina* ... 163–172
virginicus, resistance to aeciospores of *Puccinia triticina* ... 163–172
Embaphion—
concavum, resemblance to *Embaphion muricatum* ... 324
muricatum—
biology ... 323–334
control ... 333
description ... 326–332
distribution ... 324–325
enemies ... 332–333
Endothea parasitica, food of *Neoclytus capraea* ... 193
Euxoa ochragaster, connection with *Porosagrotis orthogonia* ... 290
Fasciola hepatica, hemotoxins ... 380–432
FERTILITY IN SHROPSHIRE SHEEP ... 231–234
Festuca Thurberi, host of *Puccinia Cockerelliana* ... 165
Fisher, D. F., and Brooks, Charles: TRANSPORTATION ROTS OF STONE FRUITS AS INFLUENCED BY ORCHARD SPRAYING ... 467–477
Flavone in maize ... 3–31
FLORA OF CORN MEAL ... 179–188
Fraxinus sp., host of—
Neoclytus capraea ... 210–211
Xylotrechus colonus ... 195–198
Fruit fungi, temperature relations ... 451–465
Fruits, stone, influence of orchard spraying on rots ... 467–477
Fungi, of stone fruits, temperature relations ... 451–465
FURTHER STUDIES ON RELATION OF SULPHATES TO PLANT GROWTH AND COMPOSITION ... 101–110
Fusarium—
eumartii, effect of temperature ... 66–80
hyperocysporum, effect of humidity ... 71–80
oxysporum—
cause of rot of Irish potatoes ... 65–80
confused with causal organism of black-leg potato tuber-rot ... 81–92
radicicola—
confused with causal organism of black-leg potato tuber-rot ... 81–92
effect of temperature ... 66–80

Page

Fusarium—Continued.
trichothecioides, effect of temperature ... 66–80
sp.—
in corn meal ... 187–188
on pitted grapefruit ... 277
trichothecioides, isolated from potato black-leg tubers ... 82
tuberivorum, effect of temperature ... 66–80
vasinfectum var. *pisi*, host selection ... 191–220
Gardner, Max W., and Kendrick, James B.: TURNIP MOSAIC ... 123–124
SOYBEAN MOSAIC ... 111–114
GENETIC BEHAVIOR OF THE SPELT FORM IN CROSSES BETWEEN TRITICUM SPELTA AND TRITICUM SATIVUM ... 335–364
Geographical location, effect on coniferous tree seed in storage ... 508–510
Georgia velvet bean. See *Stizolobium deeringianum*.
Glechoma hederacea, susceptibility to *Puccinia triticina* ... 152–172
Globulins in Georgia velvet bean ... 15
Glucosid of quercetin in maize ... 2–31
Gluten, effect on growing chicks ... 144
Goss, R. W.: TEMPERATURE AND HUMIDITY STUDIES OF SOME FUSARIA ROTS OF THE IRISH POTATO ... 65–80
Gracilaria stigmatella, change in nidification ... 191
Grape—
See *Vitis vinifera*.
wild. See *Vitis* sp.
Grapefruit, ripening and storage ... 263–279
Graphidium strigosum, hemotoxins ... 387–432
Grass, Sudan, hydrocyanic acid ... 125–138
Griffee, Fred: COMPARATIVE VIGOR OF F_1 WHEAT CROSSES AND THEIR PARENTS ... 53–63
Ground squirrel. See *Citellus richardsoni*.
Growth of plants, effect of sulphur ... 101–110
Hackberry. See *Celtis occidentalis*.
Haemonchus contortus, hemotoxins ... 387–432
Halbersleben, D. L., et al.: NUTRIENT REQUIREMENTS OF GROWING CHICKS; NUTRITIVE DEFICIENCIES OF CORN ... 139–149
Harlan, Harry V., and Pope, Merritt N.: ASH CONTENT OF THE AWN, RACHIS, PALEA, AND KERNEL OF BARLEY DURING GROWTH AND MATURATION ... 433–449
Harter, L. L., and Weimer, J. L.:
A COMPARISON OF THE PECTINASE PRODUCED BY DIFFERENT SPECIES OF RHIZOPUS ... 371–377
SUSCEPTIBILITY OF THE DIFFERENT VARIETIES OF SWEET POTATOES TO DECAY BY RHIZOPUS NIGRICANS AND RHIZOPUS TRITICI ... 511–515
Hawkins, Lon A.: A PHYSIOLOGICAL STUDY OF GRAPEFRUIT RIPENING AND STORAGE ... 263–279
HEMOTOXINS FROM PARASITIC WORMS ... 379–432

Page

Hepatica sp., resistance to teliospores of *Puccinia triticina* 155–172
Heterodera schachti, host selection 192
Histidin in Georgia velvet bean 15
Hogan, A. G., et al.: INFLUENCE OF THE PLANE OF NUTRITION ON THE MAINTENANCE REQUIREMENT OF CATTLE 115–121
Hoowkorm—
of cattle. See *Bustomum phlebotomum*.
of dog. See *Ancylostoma caninum*.
of man. See *Necator americanus*.
of man, Old World. See *Ancylostoma duodenale*.
HOPKINS HOST-SELECTION PRINCIPLE AS RELATED TO CERTAIN CERAMBYCID BEETLES 189–220
Hordeum—
caespitosum, resistance to aeciospores of *Puccinia triticina* 163–172
gussoneanum, resistance to aeciospores of *Puccinia triticina* 163–172
jubatum—
host of *Puccinia Agropyri* 165
resistance to aeciospores of *Puccinia triticina* 163–172
murinum, resistance to aeciospores of *Puccinia triticina* 163–172
pusillum, resistance to aeciospores of *Puccinia triticina* 163–172
vulgare, resistance to aeciospores of *Puccinia triticina* 163–172
Horned lark. See *Otocoris alpestris leucolaema*.
Host-selection principle 189–220
Hucker, G. J.: MICROSCOPIC STUDY OF BACTERIA IN CHEESE 93–100
Humidity, effect on Fusaria rots of Irish potatoes 65–80
HYDROCYANIC ACID IN SUDAN GRASS 125–138
Hydrogen-ion concentration, effect on development of nodules on legumes 30
Hydrophyllum appendiculatum, resistance to teliospores of *Puccinia triticina* 155–172
Hydroxid, sodium, effect on hydrocyanic acid in Sudan grass 135–136
Hylotrupes ligneus, host selection 207–210
Hyperplatys—
aspersus, host selection 217–219
maculatus, host selection 217–219
Hystrix Hystrix, resistance to aeciospores of *Puccinia triticina* 163–172
Impatiens sp., resistance to teliospores of *Puccinia triticina* 155–172
INFLUENCE OF THE PERIOD OF TRANSPLANTING WESTERN WHITE PINE SEEDLINGS UPON THEIR BEHAVIOR IN NURSERY AND PLANTATION 33–46
INFLUENCE OF THE PLANE OF NUTRITION ON THE MAINTENANCE REQUIREMENT OF CATTLE 115–121
Ipomoea batatas, susceptibility of different varieties to *Rhizopus* spp 511–515
Irish potato. See *Solanum tuberosum*.

Page

Iron sulphate, effect on growing chicks 145–149
Jackson, H. S., and Mains, E. B.: AECIAL STAGE OF THE ORANGE LEAF-RUST OF WHEAT, PUCCINIA TRITICINA ERIKS 151–172
Jones, Fred Reuel, and Tisdale, W. B.: EFFECT OF SOIL TEMPERATURE UPON THE DEVELOPMENT OF NODULES ON THE ROOTS OF CERTAIN LEGUMES 17–31
Jones, L. R., and Walker, J. C.: RELATION OF SOIL TEMPERATURE AND OTHER FACTORS TO ONION SMUT INFECTION 235–262
Kafir, hydrocyanic acid 137
Kendrick, James B., and Gardner, Max W.: SOYBEAN MOSAIC 111–114
TURNIP MOSAIC 123–124
Kernel of barley, ash content 433–449
Knautia arvensis, susceptibility to *Puccinia triticina* 152–172
Lactate, calcium, effect on growing chicks 145–149
Lark, horned. See *Otocoris alpestris leucolaema*.
Launaea asplenifolia, suspected aecial host of *Puccinia triticina* 153–172
Leafrust, orange. See *Puccinia triticina*.
"Leak," confused with blackeg potato tuber-rot 81–92
Lecanium—
corni, host selection 191
robiniarum, host selection 191–220
LeFevre, Edwin, and Thom, Charles: FLORA OF CORN MEAL 179–188
Legumes, effect of soil temperature on development of nodules 17–31
Leighty, Clyde E., and Boshnakian, Sarkis: GENETIC BEHAVIOR OF THE SPELT FORM IN CROSSES BETWEEN TRITICUM SPELTA AND TRITICUM SATIVUM 335–364
Ligustrum vulgare, susceptibility to *Puccinia triticina* 152–172
Liopus alpha, host selection 194–220
Liriodendron tulipifera—
host of *Hyperplatys maculatus* 217–219
immunity to *Neoclytus erythrocephalus* 213–215
Lithospermum purpureo-coeruleum, susceptibility to *Puccinia triticina* 152–172
Locust. See *Robinia pseudacacia*.
Lodgepole pine. See *Pinus contorta*.
Lythrum Salicaria, susceptibility to *Puccinia triticina* 152–172
Macracanthorhynchus hirudinaceus, hemotoxins 387–432
Magnesium sulphate, effect on growing chicks 145–149
Mains, E. B., and Jackson, H. S.: AECIAL STAGE OF THE ORANGE LEAF-RUST OF WHEAT, PUCCINIA TRITICINA ERIKS 151–172
Maize—
nutritive deficiencies 139–149
quercetin in brown-husked 1–31
Malic acid. See Acid, malic.
Maple, red. See *Acer rubrum*.

Page

Maple, sugar. See *Acer saccharum*.
Meal, corn, flora 179–188
Melandryum album, susceptibility to *Puccinia triticina* 152–172
Mertensia virginica, resistance to teliospores of *Puccinia triticina* 155–172
Mesquite. See *Prosopis juliflora*.
Metarrhizium anisophae, parasite of *Embaphion muricatum* 332
MICROSCOPIC STUDY OF BACTERIA IN CHEESE 93–100
Miller, Harry G.: FURTHER STUDIES ON RELATION OF SULPHATES TO PLANT GROWTH AND COMPOSITION 101–110
Mold, blue. See *Penicillium expansum*.
Mold rot, black, caused by *Rhizopus nigricans* 451–465
Molds in corn meal 181–188
Molorchus bimaculatus, host selection 212–213
Monilia. See *Sclerotinia cinerea*.
Morus rubra, host of *Cyllene pictus* 198–203
Mosaic—
of Chinese cabbage 173–178
of soybean 111–114
of turnip 123–124
Mulberry. See *Morus rubra*.
Multiceps multiceps, hemotoxins 382–432
Mumford, F. B., Hogan, A. G., and Salmon, W. D.: INFLUENCE OF THE PLANE OF NUTRITION ON THE MAINTENANCE REQUIREMENT OF CATTLE 115–121
Mussehl, F. E., Calvin, J. W., Halbersleben, D. L., and Sandstedt, R. M.: NUTRIENT REQUIREMENTS OF GROWING CHICKS: NUTRITIVE DEFICIENCIES OF CORN 139–149
Mustard. See *Brassica japonica*.
Myosotis—
alpestris, failure to become infected from basidiospores of *Puccinia rubigo-vera* ... 152–172
arvensis, failure to become infected from basidiospores of *Puccinia rubigo-vera* ... 152–172
palustris, resistance to teliospores of *Puccinia triticina* 155–172
scorpoides, resistance to teliospores of *Puccinia triticina* 155–172
sp., susceptibility to *Puccinia triticina* ... 152–172
Myzus persicae, carriers of mosaic of Chinese cabbage 173–178
Nasturtium sp., susceptibility to *Puccinia triticina* 152–172
Necator americanus, hemotoxins 383–432
Necosmopora vasinfecta, host selection 191–220
Neoclytus—
capraea, host selection 193–220
erythrocephalus, host selection 213–215
Nitrate, sodium, effect on plant growth 102–110
Nitrates, effect on development on nodules on legumes 18
Nitrogen, effect of soil temperature on assimilation by legumes 17–31
Nodules, effect of soil temperature on development 17–31

Page

Nonnea rosea, failure to become infected from basidiospores of *Puccinia rubigo-vera* 152–172
NOTES ON ORGANIC ACIDS OF PYRUS CORONARIA, RHUS GLABRA, AND ACER SACCHARUM 221–230
Notholcus lanatus, resistance to aeciospores of *Puccinia triticina* 163–172
NUTRIENT REQUIREMENTS OF GROWING CHICKS: NUTRITIVE DEFICIENCIES OF CORN 139–149
Oak, white. See *Quercus alba*.
OCCURRENCE OF QUERCETIN IN EMERSON'S BROWN-HUSKED TYPE OF MAIZE 1–31
(*Oceneria*) *Porthetria dispar*, host selection 191
Oncideres cingulata, host selection 194–220
Onion smut. See *Urocystis cepulae*.
Orange leafrust. See *Puccinia triticina*.
Orange, osage. See *Toxylon pomiferum*.
Ornithogalum umbellatum, resistance to teliospores of *Puccinia triticina* 155–172
Osage orange. See *Toxylon pomiferum*.
Otocoris alpestris leucolaema, enemy of *Porosogrotis orthogonia* 313
Palea of barley, ash content 433–449
PALE WESTERN CUTWORM (POROSAGROTIS ORTHOGONIS MORR.) ... 289–322
Paloverde. See *Parkinsonia microphylla*.
Parker, J. R., Strand, A. L., and Seamans, H. L.: PALE WESTERN CUTWORM (POROSAGROTIS ORTHOGONIA MORR.) 289–322
Parkinsonia microphylla, host of *Cyllene crinicornis* 203
Pastinaca sativa, susceptibility to *Puccinia triticina* 152–172
Peach. See *Amygdalus persica*.
Pectinase produced by species of Rhizopus. 371–377
Peleteria robusta, parasite of *Porosogrotis orthogonia* 313
Pemphigus betae, carrier of *corticium vagum* ... 50–52
Penicillium—
expansum in corn meal 185–188
luteum in corn meal 185–188
oxalicum in corn meal 185–188
purpurogenum 185–188
sp. on pitted grapefruit 277
Penta-acetyl-quercetin in maize 3–31
Perilitus eleodis, parasite of *Embaphion muricatum* 332
Phacelia—
Purshii, resistance to teliospores of *Puccinia triticina* 155–172
tanacetifolia, resistance to teliospores of *Puccinia triticina* 155–172
Phillyrea sp., susceptibility to *Puccinia triticina* 152–172
Phosphate, dipotassium, effect on growing chicks 145–149
Phosphoric acid. See Acid, phosphoric.
Phratora vitellinae, inherited changes in habits 191
Phyllosticta—
congesta, cause of plum blotch 365–370
solitaria, cause of apple blotch 365

Page

PHYSIOLOGICAL STUDY OF GRAPE-FRUIT RIPENING AND STORAGE, A 263-279
Picea engelmanni, storage of seed 481-510
Pigmentation of maize 1-31
Pine—
lodgepole. See *Pinus contorta*.
western white. See *Pinus monticola*.
yellow. *Pinus ponderosa*.
Pine beetle. See *Dendroctonus monticolae*.
Pinus—
contorta—
host of *Dendroctonus monticolae* 189-220
storage of seed 481-510
monticola—
influence of period of transplanting on growth of seedlings 33-46
storage of seed 481-510
ponderosa—
threatened by *Dendroctonus monticolae*. 189-220
storage of seed 481-510
strobus, storage of seed 481-510
PLUM BLOTCH, A DISEASE OF THE JAPANESE PLUM, CAUSED BY PHYLLOSTICTA CONGESTA HEALD AND WOLF 365-370
Plum, Japanese. See *Prunus triflora*.
Poa nemoralis var. *firmula*, host of *Puccinia triticina* 164-172
Poplar, yellow. See *Liriodendron tulipifera*.
Porosagrotis—
delorata. Syn. *Porosagrotis orthogonia*.
orthogonia—
control 307-313
distribution 291
economic importance 305-306
history 289-290
life history 293-305, 314-320
natural enemies 313-314
Porthetria dispar, host selection 191-220
Pope, Merritt N., and Harlan, Harry V.: ASH CONTENT OF THE AWN, RACHIS, PALEA, AND KERNEL OF BARLEY DURING GROWTH AND MATURATION 433-449
Potato—
Irish. See *Solanum tuberosum*.
sweet. See *Ipomoea batatas*.
Potato tuber-rot, blackleg 81-92
Prosopis juliflora, host of *Cyllene crinicornis* .. 203
Prunella vulgaris, susceptibility to *Puccinia triticina* 152-172
Prunes, effect of—
orchard spraying on transportation rots .. 471-477
temperature on damage from rots 452-465
Prunus triflora, host of *Phullosticta congesta* . 365-370
Pseudotsuga taxifolia, storage of seed 481-510
Puccinia—
Agropyri—
æcia on Ranunculaceous hosts 154
rust on *Thalictrum dasycarpum* 165
alternans—
æcia on Ranunculaceous hosts 154
rust on *Bromus Porteri* 165
borealis, rust on *Agrostis borealis* 164

Page

Puccinia—Continued.
bromina, pathogenicity to *Symphytum officinale* 154-172
Clematidis, æcia 151-172
Cockerelliana, rust on *Festuca Thurberi* 165
dispersa, brown rust of wheat 152-172
glumarum, on wheat 151-172
graminis, on wheat 151-172
obliterata, rust on *Thalictrum alpinum* 165
perplexans, æcia on Ranunculaceous hosts .. 154
persistens, æcia on Ranunculaceous hosts .. 154
rubigo-vera, on rye 151-172
straminis. Syn. *Puccinia rubigo-vera*.
triticina, æcial stage 151-172
Pulmonaria—
montana, host of *Puccinia bromina* 154-172
officinalis, failure to become infected from basidiospores of *Puccinia rubigo-vera* 152-172
Pyrus coronaria, organic acids 221-229
Pythium debaryanum, confused with causal organism of blackleg potato tuber-rot 81-92
Quamasia hyacinthina. Syn. *Camassia esculenta*.
Quercetin, in brown-husked maize 1-31
Quercus alba, host of *Neoclytus capraea* 210-211
Rachis of barley, ash content 433-449
Ragweed. See *Ambrosia artemisiifolia*.
Ram, relation of age to fertility 232-233
Ranunculus—
susceptibility to *Puccinia triticina* 152-172
aconitifolius, resistance to teliospores of *Puccinia triticina* 155-172
acris—
failure to become infected from basidiospores of *Puccinia rubigo-vera* 152-172
resistance to teliospores of *Puccinia triticina* 155-172
asiaticus, susceptibility to *Puccinia triticina* 152-172
auricomus, susceptibility to *Puccinia triticina* 152-172
bulbosus, failure to become infected from basidiospores of *Puccinia rubigo-vera* 152-172
Ficaria, susceptibility to *Puccinia triticina* 152-172
flammula, susceptibility to *Puccinia triticina* 152-172
lanuginosus, susceptibility to *Puccinia triticina* 152-172
repens, susceptibility to *Puccinia triticina*. 152-172
Read, J. W., and Sure, Barnett.: BIOLOGICAL ANALYSIS OF THE SEED OF THE GEORGIA VELVET BEAN, STIZOLOBIUM DEERINGIANUM 5-15
Redbud. See *Cercis canadensis*.
Red maple. See *Acer rubrum*.
RELATION OF SOIL TEMPERATURE AND OTHER FACTORS TO ONION SMUT INFECTION 235-262
Rhamus—
cathartica, failure to become infected from basidiospores of *Puccinia rubigo-vera* 152-172
Frangula, failure to become infected from basidiospores of *Puccinia rubigo-vera* 152-172
Rhizoctonia solani, similarity to *Corticium vagum* 51

Page

Rhizopus—
arrhizus—
pathogenicity on sweet potato..... 511
production of pectinase..... 372-377
artocarpi—
production on pectinase..... 372-377
pathogenicity on sweet potato..... 511
chinensis, production of pectinase..... 372-377
delemar—
pathogenicity on sweet potato..... 511
production of pectinase..... 372-377
maydis—
pathogenicity on sweet potato..... 511
production of pectinase..... 372-377
microsporus, production of pectinase..... 372-377
nigricans—
in corn meal..... 185-188
pathogenicity on different varieties of sweet potatoes..... 511-515
production of pectinase..... 372-377
temperature relations..... 451-465
nodosus—
pathogenicity on sweet potato..... 511
production of pectinase..... 372-377
oryzae—
pathogenicity on sweet potato..... 511
production of pectinase..... 372-377
reflexus—
pathogenicity on sweet potato..... 511
production of pectinase..... 372-377
spp., production of pectinase..... 371-377
tritici—
pathogenicity on sweet potato..... 511
production of pectinase..... 371-377
Rhus glabra, organic acids..... 221-229
Richards, B. L.: A DRYROT CANKER OF SUGAR BEETS..... 47-52
Ribes Grossularia, susceptibility to *Puccinia triticina*..... 152-172
Roberts, Elmer: FERTILITY IN SHROPSHIRE SHEEP..... 231-234
Roberts, John W.: PLUM BLOTCH, A DISEASE OF THE JAPANESE PLUM, CAUSED BY PHYLLOSTICTA CONGESTA HEALD AND WOLF..... 365-370
Robinia pseudacacia, host of—
Lecanium corni..... 191-220
Xylotrechus colonus..... 195-198
Rogers, E. C.: INFLUENCE OF THE PERIOD OF TRANSPLANTING WESTERN WHITE PINE SEEDLINGS UPON THEIR BEHAVIOR IN NURSERY AND PLANTATION..... 33-46
Rot—
black mold, caused by *Rhizopus nigricans* 451-465
blue mold. See *Penicillium expansum*.
brown, caused by *Sclerotinia cinerea*..... 451-465
Rots, transportation, influence of orchard spraying..... 467-477
Roughage requirements of chicks..... 146
Rumex acetosa, susceptibility to *Puccinia triticina*..... 152-172
Salmon, W. D., et al.: INFLUENCE OF THE PLANE OF NUTRITION ON THE MAINTENANCE REQUIREMENT OF CATTLE..... 115-121

Page

Sando, Charles E., and Bartlett, H. H.: NOTES ON THE ORGANIC ACIDS OF PYRUS CORONARIA, RHUS GLABRA, AND ACER SACCHARUM.. 221-229
OCCURRENCE OF QUERCETIN IN EMERSON'S BROWN-HUSKED TYPE OF MAIZE..... 1-31
Sandstedt, R. M., et al.: NUTRIENT REQUIREMENTS OF GROWING CHICKS: NUTRITIVE DEFICIENCIES OF CORN..... 139-149
Schistosoma japonicum, hemotoxins..... 387-432
Schultz, E. S.: A TRANSMISSIBLE MOSAIC DISEASE OF CHINESE CABBAGE, MUSTARD, AND TURNIP..... 173-178
Schwartz, Benjamin: HEMOTOXINS FROM PARASITIC WORMS..... 379-432
Sclerotinia cinerea, temperature relations... 451-465
Seamans, H. L., et al.: PALE WESTERN CUTWORM (POROSAGROTIS ORTHOGONIA MORR.)..... 289-322
Secale cereale, susceptibility to aeciospores of *Puccinia triticina*..... 163-172
Seed of coniferous trees, storage..... 479-510
Sequoia sempervirens, host of *Hylotrupes ligneus*..... 209-210
Serum, effect on hookworm hemolysin..... 412-413
Shapovalov, M., and Edson, H. A.: BLACK-LEG POTATO TUBER-ROT UNDER IRRIGATION..... 81-92
Sheep, Shropshire, fertility..... 231-234
Shropshire sheep, fertility..... 231-234
Sitanion Hystrix, susceptibility to aeciospores of *Puccinia triticina*..... 163-172
Smut—
of oats. See *Ustilago avenae*.
onion. See *Urocystis cepulae*.
stinking, of wheat. See *Tilletia tritici*.
Sodium—
carbonate, effect on hydrocyanic acid in Sudan grass..... 135-136
chlorid, effect on growing chicks..... 145-149
hydroxid, effect on hydrocyanic acid in Sudan grass..... 135-136
nitrate, effect on plant growth..... 102-110
sulphate, effect on plant growth..... 102-110
Soil temperature, effect on—
nodules..... 17-31
onion smut..... 235-262
Solanum tuberosum—
absorption of copper from soil..... 281-287
Fusaria rots..... 65-80
Sorghums, hydrocyanic acid..... 125-138
Soybean meal, effect on growing chicks..... 145
SOYBEAN MOSAIC..... 111-114
Soybeans, effect of soil temperature on nodule development..... 23
Sparrow, western grasshopper. See *Ammodramus savannarum bimaculatus*.
Spelt form, behavior in wheat crosses..... 335-364
Spraying, influence on transportation rots of stone fruits..... 467-477
Squirrel, ground. *Citellus richardsoni*.
Sterility in wheat crosses..... 58-61

Page

Stomach worm of sheep. See *Haemonchus contortus*.
Stone fruit fungi, temperature relations.... 451-465
Stone fruits, influence of orchard spraying on rots..... 467-477
STORAGE OF CONIFEROUS TREE SEED..... 479-510
Strand, A. L., et al.: PALE WESTERN CUTWORM (POROSAGROTIS ORTHOGONIA MORR.)..... 289-322
Stinking smut of wheat. See *Tilletia tritici*.
Succinic acid. See Acid, succinic.
Sudan grass. See Grass, Sudan.
Sugar-beet root aphis. See *Pemphigus betae*.
Sugar beets, dry-rot canker..... 47-52
Sugar maple. See *Acer saccharum*.
Sugars in grapefruit..... 263-279
Sulphate—
calcium, effect on plant growth..... 102-110
copper, toxicity to potato plants..... 281-287
iron, effect on growing chicks..... 145-149
magnesium, effect on growing chicks..... 139-149
sodium, effect on plant growth..... 102-110
Sulphates, relation to plant growth..... 101-110
Sulphur, effect on growing chicks..... 145-149
Sulphuric acid. See Acid, sulphuric.
Sumac. See *Rhus glabra*.
Sure, Barnett, and Read, J. W.: BIOLOGICAL ANALYSIS OF THE SEED OF THE GEORGIA VELVET BEAN, STIZOLOBIUM DEERINGIANUM.... 5-15
SUSCEPTIBILITY OF THE DIFFERENT VARIETIES OF SWEET POTATOES TO DECAY BY RHIZOPUS NIGRICANS AND RHIZOPUS TRITICI. 511-515
Swanson, C. O.: HYDROCYANIC ACID IN SUDAN GRASS..... 125-138
Sweet potatoes. See *Ipomoea batatas*.
Symphytum—
asperrimum, failure to become infected from basidiospores of *Puccinia rubigo-vera*..... 152-172
officinale—
host of *Puccinia bromina*..... 154-172
susceptibility to *Puccinia triticina*..... 152-172
Taenia—
saginata, hemotoxins..... 380-432
solium, hemotoxins..... 380, 432
Tapeworm. See *Diphyllobothrium latum*.
Taraxacum officinalis, failure to become infected from basidiospores of *Puccinia rubigo-vera*..... 152-172
TEMPERATURE AND HUMIDITY STUDIES OF SOME FUSARIA ROTS OF THE IRISH POTATO..... 65-80
Temperature—
effect on hookworm hemolysin..... 413-414
soil, effect on—
nodules..... 17-31
onion smut..... 235-262
TEMPERATURE RELATIONS OF STONE FRUIT FUNGI..... 451-465
Thalictrum—
anqustifolium, susceptibility to teliospores of *Puccinia triticina*..... 155-172

Page

Thalictrum—Continued.
aquilegifolium, susceptibility to teliospores of *Puccinia triticina*..... 155-172
dasycarpum, susceptibility to teliospores of *Puccinia triticina*..... 157-172
Delavayi, susceptibility to teliospores of *Puccinia triticina*..... 157-172
dioicum, resistance to teliospores of *Puccinia triticina*..... 155-172
Fendleri, host of *Puccinia Cockerelliana*.... 165
foetidum, host of *Puccinia triticina*..... 164-172
minus—
resistance to teliospores of *Puccinia triticina*..... 155-172
adiantifolium, susceptibility to teliospores of *Puccinia triticina*..... 157-172
polycarpum, susceptibility to teliopores of *Puccinia triticina*..... 157-172
polygamum, resistance to teliospores of *Puccinia triticina*..... 155-172
sparsiflorum, host of *Puccinia alternans*.... 165
Thom, Charles, and LeFevre, Edwin: FLORA OF CORN MEAL..... 179-188
Thysanosoma actinioides, hemotoxins..... 421-432
Tilletia tritici, effect of—
soil moisture..... 240
soil temperature..... 242
Tillotson, C. R.: STORAGE OF CONIFEROUS TREE SEED..... 479-510
Tisdale, W. B., and Jones, Fred Reuel: EFFECT OF SOIL TEMPERATURE UPON THE DEVELOPMENT OF NODULES ON THE ROOTS OF CERTAIN LEGUMES..... 17-31
Toxylon pomiferum, host of *Cyllene pictus*.. 198-203
TRANSMISSIBLE MOSAIC DISEASE OF CHINESE CABBAGE, MUSTARD, AND TURNIP, A..... 173-178
TRANSPORTATION ROTS OF STONE FRUITS AS INFLUENCED BY ORCHARD SPRAYING..... 467-477
Trichostrongylus retortaeformis, hemotoxins. 387-432
Trichuris—
depressiuscula, hemotoxins..... 420
triciura, hemotoxins..... 384-432
Triticum—
aegilops, susceptibility to aeciospores of *Puccinia triticina*..... 163-172
aestivum, host of *Puccinia triticina*..... 151-172
capitatum, behavior of the spelt form in crosses..... 335-364
compactum—
behavior of the spelt form in crosses... 335-364
vigor of F_1 of crosses..... 54-63
dicoccum—
behavior of spelt form in crosses..... 335
vigor of F_1 of crosses..... 54-63
dicoccoides, behavior of spelt form in crosses 361
durum—
behavior of spelt form in crosses..... 335
vigor of F_1 of crosses..... 54-63
polonicum, effect of crossing..... 335-364
sativum, behavior of spelt form in crosses with *T. spelta*..... 335-346

Page

Triticum—Continued.
spelta, behavior of spelt form in crosses with *T. sativum* 335-364
turgidum, behavior of spelt form in crosses. 335
vulgare—
behavior of spelt form in crosses 335-364
vigor of F_1 of crosses 54-63
Syn. *Triticum aestivum*.
Trollius europeus, resistance to teliospores of *Puccinia triticina* 155-172
Tuber-rot, blackleg 81-9
Tulip tree. See *Liriodendron tulipifera*.
Turnip. See *Brassica rapa*.
TURNIP MOSAIC 123-12
Tussilago Farfara, susceptibility to *Puccinia triticina* 152-172
Urocystis cepulae, effect of soil temperature 235-262
Urtica dioica—
failure to become infected from basidiospores of *Puccinia rubigo-vera* 152-172
susceptibility to *Puccinia triticina* 152-172
Ustilago—
avenae, effect of soil temperature 242
levis, effect of soil temperature 242
Valeriana dioica, susceptibility to *Puccinia triticina* 152-172
Velvet bean. See *Stizolobium deeringianum*.
Vitis—
vinifera, succinic acid 224
sp., host of *Cyllene pictus* 198-203

Page

Wade, J. S., and Boving, Adam G.: BIOLOGY OF EMBAPHION MURICATUM 323-334
Walker, J. C., and Jones, L. R.: RELATION OF SOIL TEMPERATURE AND OTHER FACTORS TO ONION SMUT INFECTION 235-262
Water, hot, effect on hydrocyanic acid in Sudan grass 131-133
Weather, effect on hydrocyanic acid in Sudan grass 136
Weimer, J. L., and Harter, L. L.:
A COMPARISON OF THE PECTINASE PRODUCED BY DIFFERENT SPECIES OF RHIZOPUS 371-377
SUSCEPTIBILITY OF THE DIFFERENT VARIETIES OF SWEET POTATOES TO DECAY BY RHIZOPUS NIGRICANS AND RHIZOPUS TRITICI 511-515
Western white pine. See *Pinus monticola*.
Wheat crosses, vigor of F_1 seedlings 53-63
Wheat, orange leafrust 151-172
Whipworm. See *Trichuris trichiura*.
White oak. See *Quercus alba*.
White pine, western. See *Pinus monticola*.
Worms, parasitic, homotoxins 379-432
Xylotrechus colonus, host selection 195-220
Yellow pine. See *Pinus ponderosa*.
Yellow poplar. See *Liriodendron tulipifera*.